

New chiral phases of superfluid ^3He stabilized by anisotropic silica aerogel

J. Pollanen, J. I. A. Li, C. A. Collett, W. J. Gannon, W. P. Halperin* and J. A. Sauls

A rich variety of Fermi systems condense by forming bound pairs, including high-temperature¹ and heavy-fermion² superconductors, Sr_2RuO_4 (ref. 3), cold atomic gases⁴ and superfluid ^3He (ref. 5). Some of these form exotic quantum states with non-zero orbital angular momentum. We have discovered, in the case of ^3He , that anisotropic disorder, engineered from highly porous silica aerogel, stabilizes a chiral superfluid state that otherwise would not exist. Furthermore, we find that the chiral axis of this state can be uniquely oriented with the application of a magnetic field perpendicular to the aerogel anisotropy axis. At sufficiently low temperature we observe a sharp transition from a uniformly oriented chiral state to a disordered structure consistent with locally ordered domains, contrary to expectations for a superfluid glass phase⁶.

Superconducting states with non-zero orbital angular momentum, $L \neq 0$, are characterized by a competitive, but essential, relationship with magnetism, strong normal-state anisotropy or both^{1–3,5}. Moreover, these states are strongly suppressed by disorder, an important consideration for applications⁷ and a signature of their unconventional behaviour^{3,8,9}. Although liquid ^3He in its normal phase is perfectly isotropic, it becomes a p -wave superfluid at low temperatures with non-zero orbital and spin angular momenta, $L = S = 1$ (ref. 10). One of its two superfluid phases in zero magnetic field is anisotropic with chiral symmetry, where the handedness results from the orbital motion of the bound ^3He pairs about an axis ℓ . This chiral superfluid, called the A phase or axial state, is stable at high pressure near the normal-to-superfluid transition, Fig. 1a–c, whereas the majority of the phase diagram is the non-chiral B phase, with isotropic physical properties. The stability of the A phase is attributed to strong-coupling effects arising from collisions between ^3He quasiparticles¹⁰. However, in the presence of isotropic disorder these strong-coupling effects are reduced and the stable chiral phase disappears^{11,12}, Fig. 1a. Here we show that anisotropic disorder can reverse this process and stabilize an anisotropic phase over the entire phase diagram, Fig. 1c.

For many years it was thought to be impossible to introduce disorder into liquid ^3He because it is intrinsically chemically and isotopically pure at low temperatures. Then it was discovered^{13,14} that ^3He imbedded in $\sim 98\%$ porosity silica aerogel, Fig. 1d, is a superfluid with a transition temperature that is sharply defined¹², but reduced from that of pure ^3He . To test predictions that isotropic disorder favours isotropic states and anisotropic disorder favours anisotropic states¹⁵, we have grown a 97.5% porosity anisotropic aerogel with growth-induced radial compression¹⁶, effectively stretching it along its cylinder axis by 14.3%. Experiments using uncharacterized stretched aerogels have been previously reported^{17,18} and are in disagreement with the work presented here. Silica aerogels, as in Fig. 1d, are formed by silica particles ≈ 3 nm in diameter, precipitated from a tetramethylorthosilicate solution, and aggregated in a diffusion-limited process. After supercritical drying we obtain a

cylinder as shown in Fig. 1d. Our numerical simulation of this process indicates that the gel particles form strands (Fig. 1d). According to theory¹⁹, ℓ is constrained to point outward from the strand. Stretching an aerogel tends to align the strands, thereby forcing the chiral axis, on average, to be in an easy plane perpendicular to the aerogel anisotropy axis⁶. On the other hand, for an axially compressed aerogel ℓ should be aligned with the compression axis⁶.

We use pulsed nuclear magnetic resonance (NMR; Methods), to identify the ^3He superfluid state and determine the magnitude of its order parameter, Δ . The equilibrium nuclear magnetization is tipped by a pulsed magnetic field to an angle β away from the static applied magnetic field, \mathbf{H} , and the Fourier transform of its free precession is the NMR spectrum. The frequency shift of the spectrum, $\Delta\omega$, from the normal-state value centred at the Larmor frequency, ω_L , is a direct measure of the temperature-dependent order parameter, $\Delta(T)$. For the A and B phases of ^3He (ref. 10) the frequency shifts are

$$\text{A phase: } \Delta\omega_A(\beta) = \frac{\Omega_A^2}{2\omega_L} \left(\frac{1}{4} + \frac{3}{4} \cos(\beta) \right); \quad \Omega_A^2 = \Delta_A^2 \left(\frac{12}{5} \frac{\gamma^2 \lambda_D N_F}{\chi_A} \right) \quad (1)$$

$$\text{B phase: } \Delta\omega_B(\beta) = 0 \quad \beta < 104^\circ; \quad \Omega_B^2 = \Delta_B^2 \left(6 \frac{\gamma^2 \lambda_D N_F}{\chi_B} \right) \quad (2)$$

where $\Omega_{A,B}^2$ is the square of the longitudinal resonance frequency, which is proportional to $\Delta_{A,B}^2$ and determined by constants of the normal fluid: gyromagnetic ratio, γ ; dimensionless dipole coupling constant, λ_D ; and single-spin density of states at the Fermi level, N_F ; as well as the magnetic susceptibilities χ_A and χ_B . The A-phase susceptibility is temperature independent and equal to the normal-state value, χ_N ; however, χ_B is temperature dependent and suppressed relative to χ_N .

For ^3He in our anisotropic aerogel we observe a sharp transition to a superfluid state at a temperature, T_{ca} , marked by the onset of NMR frequency shifts for small tip angle, $\beta = 8^\circ$ (Fig. 2a). Our comparison of the data with either of the two pure ^3He states based on equations (1) and (2) indicates that the superfluid is an axial p -wave state similarly to the A phase. In the Ginzburg–Landau (GL) limit, $T \lesssim T_c$, the initial slope of $\Delta\omega_A$ is proportional to the square of the order parameter magnitude Δ_{A0}^2 , where $\Delta_A^2 = \Delta_{A0}^2 (1 - T/T_c)$. We find in our anisotropic aerogel that Δ_{A0}^2 has a linear pressure dependence, as shown in Fig. 2b, just as is the case for pure ^3He -A (refs 20,21). It is, however, substantially reduced in magnitude and extrapolates to a critical pressure of $P_c = 7.9$ bar at $T = 0$, where the order parameter vanishes at a quantum critical point²². We have analysed the transition temperatures $T_{ca}(P)$ using GL theory²³ (Supplementary Information) to determine the mean free path $\lambda = 113$ nm and the silica particle correlation length $\xi_a = 39$ nm. Our calculated phase diagram compares very well with the data in Fig. 1c. With the same parameters we have also calculated Δ_{A0}^2 (Supplementary Information), shown in Fig. 2b, and find excellent




Figure 1 | Superfluid phase diagrams. **a–c**, Comparison of pressure–temperature (P – T) phase diagrams of ^3He in silica aerogels with various types of anisotropy in the limit $H \rightarrow 0$. High spatial uniformity of the aerogels, better than 99.97%, was established using optical cross-polarization methods¹⁶ for **a** and **c**. The solid black curves are fits to $T_{ca}(P)$ using the GL-scattering theory (Supplementary Information) and the solid (dashed) red curves correspond to the superfluid transition T_c ($B \rightarrow A$ transition T_{BA}) for pure ^3He at $H = 0$. For subscripts, the lowercase a signifies aerogel and uppercase A and B represent the corresponding phases of superfluid ^3He . **a**, A uniformly isotropic 98.2% porosity aerogel¹². **b**, A 98.0% porosity aerogel axially compressed by 10% from torsional oscillator experiments²⁵. In both **a** and **b** the B phase occupies the majority of the P – T plane. **c**, For the 97.5% porosity stretched aerogel used in the present work the A phase is stable throughout the phase diagram. The open squares (T_d) mark the transition to a disordered chiral phase and the open circle at $T = 0$ indicates a quantum critical point at which the superfluid order parameter vanishes (Fig. 2b). **d**, Left: Photograph of an aerogel with $\sim 98\%$ porosity, 1 cm in diameter, partially extracted from the glass tube in which it was grown. Centre: Perspective view of the microstructure of the aerogel based on numerical simulation of diffusion-limited aggregation of silica particles, with diameter δ , into a network of strands with most probable spacing ξ_a . Right: The enlarged sketch shows the favourable directions (red arrows) of the chiral axes, ℓ , perpendicular to the strand.

agreement with the order parameter determined from the frequency shifts, equation (1), consistent with a suppressed A phase. Moreover, extrapolation of $\Delta\omega(P) \rightarrow 0$ is at a pressure that coincides with the critical pressure from the calculated phase diagram, demonstrating important consistency between experiment and theory.

The tip-angle dependence of the NMR frequency shift is a fingerprint of a specific p -wave state¹⁰. Our measurements of $\Delta\omega(\beta)$, Fig. 2c, follow the expected behaviour of the chiral axial state²⁴, equation (1). The magnitude of the shift is reduced when compared with that of pure ^3He -A, as shown in Fig. 2b. Furthermore, the axial state belongs to the class of equal-spin-pairing (ESP) states, with the same susceptibility as the normal Fermi liquid. We have directly demonstrated that this is the case as shown in Fig. 2d.

Last, we have looked for a possible phase transition to some other p -wave state by cooling to our lowest temperatures, $\lesssim 650\ \mu\text{K}$, in a substantial magnetic field of 196 mT, and measuring χ and $\Delta\omega$ on warming. For example, if there had been a transition to a B phase on cooling we would have observed a discontinuous increase of χ on warming from the B phase to a magnetic field-induced A phase, as is characteristic of the first-order B-to-A transition in an isotropic aerogel (Fig. 2d; refs 12). Correspondingly, we would also have observed a discontinuous change in $\Delta\omega$. The absence of these discontinuities demonstrates that an ESP superfluid state in a stretched aerogel is stable down to our lowest temperatures.

Our three independent NMR measurements show that an anisotropic stretched aerogel stabilizes anisotropic superfluidity throughout the phase diagram, Fig. 1c. This is in stark contrast to ^3He in the presence of isotropic disorder^{11,12}, Fig. 1a, or for anisotropic disorder established by axial compression, Fig. 1b (ref. 25). In both of these latter cases the B phase is dominant.

Understanding the stability of chiral superfluidity, stabilized by anisotropy, is an interesting open problem, in part because topologically non-trivial pairing states have unusual structure.

Our NMR results reported in Fig. 2a–c were restricted to temperatures within $\sim 20\%$ of T_{ca} . In this high-temperature region there is negligible increase in the NMR linewidth as compared with the normal fluid (Fig. 3), indicating that the chiral axis, ℓ , has a well-defined direction orthogonal to both the aerogel anisotropy axis⁶ and the magnetic field. This orientation minimizes the magnetic dipole energy, producing the so-called dipole-locked condition, on which the validity of equation (1) depends¹⁰, and with which our tip-angle results agree. The combined effects of the dipole energy and aerogel anisotropy define a uniform direction of ℓ for the A phase of superfluid ^3He in our stretched aerogel.

At lower temperatures, we have discovered a different phase characterized by a significantly broadened NMR line. The data indicate that this superfluid state is no longer the homogeneous A phase with $\ell \perp \mathbf{H}$. On warming there is an abrupt increase in the first moment of the NMR spectrum (frequency shift) and a corresponding decrease in the linewidth at a well-defined temperature, T_d (Fig. 3). The transition at T_d is independent of magnetic field for $31.1 < H < 196$ mT. In addition, the magnetic susceptibility does not change at this temperature; however, T_d is pressure dependent, as shown by the open squares in the phase diagram of Fig. 1c. We interpret these results as evidence for a disordered ESP phase with a possible phase transition at T_d .

The frequency shift is a measure of Δ_A , given in equation (1) for the dipole-locked case where the angle between the chiral axis and the magnetic field is $\theta = \cos^{-1}(\ell \cdot \mathbf{H}) = 90^\circ$. However, in general the shift also depends on θ , where for small tip angles




Figure 2 | Identification of the superfluid state. **a**, The frequency shift of the NMR spectrum for a small tip angle, $\beta = 8^\circ$, as determined from Gaussian fits. The superfluid transition temperature at different pressures is marked by the onset of positive frequency shifts. The NMR spectra have been corrected for fast exchange of liquid spins with several layers of paramagnetic solid ^3He on the aerogel surface^{14,28}. **b**, The square of the order-parameter magnitude as a function of pressure in our aerogel (closed green circles) was calculated from the initial slopes in **a** using equation (1), showing the suppression relative to pure ^3He -A (open and closed squares; refs 20,21). The experimental results are compared with the predictions of the GL theory (open blue circles; Supplementary Information). The critical pressure, $P_c = 7.9$ bar, is determined from an extrapolation (dashed line) of Δ_{A0}^2 to zero, indicated by the arrow. The solid line is a linear fit to the pure A-phase data. **c**, Identification of the superfluid as the axial p -wave state follows from the agreement with theory²⁴ (dashed blue curve) for the dependence of frequency shift $\Delta\omega$ on tip angle, $T_d < T < T_c$. **d**, The susceptibility for stretched and isotropic aerogel samples taken on warming at a pressure $P \approx 26$ bar in $H = 196$ mT after subtraction of a paramagnetic background from solid ^3He (ref. 14). For the isotropic aerogel the field-induced A-phase region has a constant susceptibility below T_{ca} and a very sharp jump at the BA transition¹². A similar jump in susceptibility would be anticipated for the stretched aerogel if the ^3He were in the B phase or any other non-ESP phase.

$\Delta\omega(\theta)$ reduces to²⁶

$$\Delta\omega(\theta) = -\frac{\Omega_A^2}{2\omega_L} \cos(2\theta) \quad (3)$$

This enables us to extract the angular distribution, $P(\theta)$, of ℓ relative to \mathbf{H} , from the NMR spectra both above and below T_d . The observed spectrum is given by the convolution product of the normal-state lineshape and the frequency distribution $P(\omega)$ that corresponds to $P(\theta)$ determined by equation (3) (Supplementary Information). Above T_d the NMR spectrum (Fig. 4a) can be fitted with an ℓ distribution (Fig. 4b) that has a single component centred at $\theta = 90^\circ$, as sketched in Fig. 4c, with angular spread $\Delta\theta = 17.8^\circ \pm 3.6^\circ$. Examination of the spectrum below T_d , Fig. 4a, indicates that it is bimodal, suggesting that $P(\theta)$ has three main components, which we take to be Gaussian functions with adjustable position, width and relative weight. From our fits we determine these parameters and obtain $P(\theta)$, Fig. 4b. For the spectra below T_d , the disordered state is composed of three domains. Approximately one-third of the sample remains with $\ell \perp \mathbf{H}$ whereas the majority, $\sim 2/3$ of the distribution, has nearly zero frequency shift with $\theta = 43.8^\circ \pm 0.1^\circ$. The theory, equation (3), does not distinguish between angles that are symmetric about $\theta = 90^\circ$, and thus we assume that $P(\theta) = P(180^\circ - \theta)$ as shown in Fig. 4d. Our interpretation of the disordered state in terms of distinct ℓ domains is different from the two-dimensional ‘orbital glass’ phase defined by a random distribution of ℓ confined to a

plane perpendicular to the stretching axis⁶. It is noteworthy that the possible existence of orbital domains, such as we observe here, has been central to the discussion of time-reversal symmetry breaking in the unconventional superconducting state in Sr_2RuO_4 (ref. 27).

Methods

In our experiments the external magnetic field \mathbf{H} was 31.1, 95.5 or 196 mT and oriented perpendicular to the aerogel cylinder axis. The sample was 4.93 mm long and had a diameter of 3.43 mm. The susceptibility was obtained by numerically integrating the phase-corrected absorption spectrum, whereas the frequency shift and linewidth were determined from the power spectrum. The sample was cooled using adiabatic nuclear demagnetization of PrNi_5 . NMR on ^{195}Pt was used for thermometry at $H = 95.5$ and 196 mT and calibrated relative to the known phase diagram of pure superfluid ^3He from a volume of liquid outside the aerogel sample equal to 30% of the total liquid. For data acquired at $H = 31.1$ mT, temperature was determined from the Curie–Weiss dependence of the solid ^3He adsorbed to the aerogel surface^{14,28}.

Received 25 November 2011; accepted 6 January 2012;
published online 12 February 2012

References

- Kirtley, J. R. *et al.* Symmetry of the order parameter in the high- T_c superconductor $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$. *Nature* **373**, 225–228 (1995).
- Heffner, R. H. & Norman, M. R. Heavy fermion superconductivity. *Comments Condens. Matter Phys.* **17**, 361–408 (1996).
- Mackenzie, A. P. & Maeno, Y. The superconductivity of Sr_2RuO_4 and the physics of spin-triplet pairing. *Rev. Mod. Phys.* **75**, 657–712 (2005).




Figure 3 | The disorder transition. Below the disorder transition, $T < T_d$ (vertical dashed lines), the NMR frequency shift drops and the linewidth abruptly increases. The frequency shift is defined as the first moment of the NMR spectrum and the linewidth is the full-width at half-maximum. Both quantities have been corrected for the effect of fast exchange between liquid and surface solid ^3He spins^{14,28}. Data were taken on warming.

- Chin, C. *et al.* Observation of the pairing gap in a strongly interacting Fermi gas. *Science* **305**, 1128–1130 (2004).
- Osheroff, D. D., Richardson, R. C. & Lee, D. M. Evidence for a new phase of solid He^3 . *Phys. Rev. Lett.* **28**, 885–888 (1972).
- Volovik, G. E. On Larkin–Imry–Ma state of ^3He -A in aerogel. *J. Low Temp. Phys.* **150**, 453–463 (2008).
- Scanlan, R. M., Malozemoff, A. P. & Larbaestier, D. C. Superconducting materials for large scale applications. *Proc. IEEE* **92**, 1639–1654 (2004).
- Tsuneto, T. *Univ. Tokyo Tech. Rep.* Vol. 47 (Inst. Solid State Phys. Ser. A, 1962).
- Dalichaouch, Y. *et al.* Impurity scattering and triplet superconductivity in UPt_3 . *Phys. Rev. Lett.* **75**, 3938–3941 (1995).
- Vollhardt, D. & Wölfle, P. *The Superfluid Phases of Helium 3* (Taylor and Francis, 1990).
- Moon, B. H. *et al.* Ultrasound attenuation and a P – B – T phase diagram of superfluid ^3He in 98% aerogel. *Phys. Rev. B* **81**, 134526 (2010).
- Pollanen, J., Li, J. I. A., Collett, C. A., Gannon, W. J. & Halperin, W. P. Identification of superfluid phases of ^3He in uniformly isotropic 98.2% aerogel. *Phys. Rev. Lett.* **107**, 195301 (2011).
- Porto, J. V. & Parpia, J. M. Superfluid ^3He in aerogel. *Phys. Rev. Lett.* **74**, 4667–4670 (1995).
- Sprague, D. T. *et al.* Homogeneous equal-spin pairing superfluid state of ^3He in aerogel. *Phys. Rev. Lett.* **75**, 661–664 (1995).
- Thuneberg, E. V., Yip, S. K., Fogelström, M. & Sauls, J. A. Models for superfluid ^3He in aerogel. *Phys. Rev. Lett.* **80**, 2861–2864 (1998).
- Pollanen, J. *et al.* Globally anisotropic high porosity silica aerogels. *J. Non-Crystal. Solids* **354**, 4668–4674 (2008).
- Elbs, J., Bunkov, Y. M., Collin, E. & Godfrin, H. Strong orientational effect of stretched aerogel on the ^3He order parameter. *Phys. Rev. Lett.* **100**, 215304 (2008).
- Dmitriev, V. V. *et al.* Orbital glass and spin glass states of ^3He -A in aerogel. *JETP Lett.* **91**, 599–606 (2010).
- Rainer, D. & Vuorio, M. Small objects in superfluid ^3He . *J. Phys. C* **10**, 3093–3106 (1977).
- Schiffer, P. E. *Studies of the Superfluid Phases of Helium Three and The Magnetization of Thin Solid Films of Helium Three* Ph.D. thesis, Stanford Univ. (1993).
- Rand, M. R. *Nonlinear Spin Dynamics and Magnetic Field Distortion of the Superfluid ^3He -B Order Parameter*. Ph.D. thesis, Northwestern Univ. (1996).
- Matsumoto, K. *et al.* Quantum phase transition of ^3He in aerogel at a nonzero pressure. *Phys. Rev. Lett.* **79**, 253–256 (1997).




Figure 4 | Distribution of the direction of the chiral axis. **a**, The convolution product of the normal-state NMR line (red curve) with a best-fit angular distribution of the chiral axis, $P(\theta)$, gives an excellent representation (dashed black curves) of the NMR spectra for the superfluid state above (left panel, bold green curve) and below (right panel, bold green curve) the disorder transition (Supplementary Information). The spectra were obtained with $H = 31.1$ mT. **b**, The angular distributions for the chiral axis, ℓ , determined from **a**. The dashed pink curve is the distribution for $T_d < T < T_{ca}$. **c**, Stretching the aerogel along the cylinder axis forces ℓ into an easy plane perpendicular to the direction of strain⁶. To minimize the dipole energy ℓ must be perpendicular to \mathbf{H} , as illustrated. **d**, According to our analysis the distribution splits into three ℓ domains for $T < T_d$.

- Sauls, J. A. & Sharma, P. Impurity effects on the A_1 – A_2 splitting of superfluid ^3He in aerogel. *Phys. Rev. B* **68**, 224502 (2003).
- Brinkman, W. F. & Smith, H. Frequency shifts in pulsed NMR for $^3\text{He}(A)$. *Phys. Lett.* **51**, 449–450 (1975).
- Bennett, R. G. *et al.* Modification of the ^3He phase diagram by anisotropic disorder. *Phys. Rev. Lett.* **107**, 235504 (2011).
- Bunkov, Y. M. & Volovik, G. E. On the possibility of the homogeneously precessing domain in bulk ^3He -A. *Europhys. Lett.* **21**, 837–843 (1993).
- Ferguson, D. G. & Goldbart, P. M. Penetration of nonintegral magnetic flux through a domain-wall bend in time-reversal symmetry broken superconductors. *Phys. Rev. B* **84**, 014523 (2011).
- Collin, E., Triqueneaux, S., Bunkov, Y. M. & Godfrin, H. Fast-exchange model visualized with ^3He confined in aerogel: A Fermi liquid in contact with a ferromagnetic solid. *Phys. Rev. B* **80**, 094422 (2009).

Acknowledgements

We are grateful to J. M. Parpia, V. V. Dmitriev, G. E. Volovik, N. Mulders, K. R. Shirer, A. M. Mounce and Y. Lee for discussion and to the National Science Foundation, DMR-1103625, DMR-0805277 and DMR-1106315, for support.

Author contributions

Experimental work and analysis was principally carried out by J.P. assisted by J.I.A.L. with further support from C.A.C. and W.J.G. Advice and assistance was provided by W.P.H. (experiment) and J.A.S. (theory).

Additional information

The authors declare no competing financial interests. Supplementary information accompanies this paper on www.nature.com/naturephysics. Reprints and permissions information is available online at www.nature.com/reprints. Correspondence and requests for materials should be addressed to W.P.H.